



California EMS System Core Quality Measures

**Emergency Medical Services Authority
California Health and Human Services Agency**

EMSA #166 - Appendix E
EMS System Quality Improvement Program Guidelines





**HOWARD BACKER, MD, MPH, FACEP
DIRECTOR**

**DANIEL R. SMILEY
CHIEF DEPUTY DIRECTOR**

**TOM M^CGINNIS
CHIEF, EMS SYSTEMS DIVISION**

EMSA #166 – Appendix E
Released – January 2013

Funded by
California HealthCare Foundation
Project #16933
April 1, 2012- April 30, 2013

Acknowledgements

California Emergency Medical Services Authority Staff

Howard Backer, MD, MPH, FACEP, Director
Daniel Smiley, Chief Deputy Director
Tom McGinnis, Division Chief, EMS
Teri Harness, Assistant Division Chief, EMS
Hailey Pate, Data Program Analyst, EMS
Adam Davis, Staff Analyst, EMS

Task Force Members

Joe Barger MD, Medical Director, Contra Costa County EMS
David Chang PhD MBA, Director of Outcomes Research, Department of Surgery, UC
San Diego School of Medicine
Cathy Chidester RN, Director, Los Angeles County EMS
Craig Stroup, Quality Improvement Coordinator, Contra Costa County EMS
Karl Sporer MD, Medical Director, Alameda County EMS
Jan Ogar RN, Assistant Director, San Mateo County EMS
Laura Wallin RN, Quality Improvement Coordinator, Riverside County EMS
Dana Solomon, Program Director, California Ambulance Association
Dennis Carter, Clinical Education Services Coordinator, American Medical Response
Ric Maloney RN, Quality Improvement Coordinator, Sacramento Metro Fire
Jason Vega, Santa Clara County EMS
Susan Mori RN, Quality Improvement coordinator, Los Angeles County EMS

Table of Contents

EMS System Core Quality Measures Project

Statutory Authority	1
Project History	2
Introduction to Core Measures	3
Chart of Core Measures for California	5

Core Measures Specification Sheets

TRA-1 Scene time for severely injured trauma patients	8
TRA-2 Direct transport to trauma center for severely injured trauma patients meeting criteria	11
ACS-1 Aspirin administration for chest pain/discomfort	15
ACS-2 12 lead EKG performance	18
ACS-3 Scene time for suspected heart attack patients	20
ACS-4 Advance hospital notification for acute coronary syndrome	22
ACS-5 Direct transport to PCI center for suspected ACS patients meeting criteria	24
CAR-1 AED prior to EMS arrival	26
CAR-2 Out-of-hospital cardiac arrests return of spontaneous circulation	28
CAR-3 Out-of-hospital cardiac arrests survival to emergency department discharge	30
CAR-4 Out-of-hospital cardiac arrests survival to hospital discharge	32
STR-1 Identification of stroke in the field	34
STR-2 Glucose testing for suspected	36
STR-3 Scene time for suspected stroke patients	38
STR-4 Advance hospital notification for suspected stroke	40
STR-5 Direct transport to stroke center for suspected stroke patients meeting criteria	42
RES-1 CPAP given for patients with respiratory distress	44
RES-2 Beta2 agonist administration	46
PED-1 Pediatric asthma patients receiving bronchodilators	49
PED-2 Transport to pediatric trauma center	52
PAI-1 Pain intervention	56
PAI-2 Results of pain intervention	58
SKL-1 Endotracheal intubation success rate.....	61
SKL-2 End-tidal CO2 performed on any endotracheal intubation	63
RST-1 Ambulance response time by ambulance zone (Emergency).....	65
RST-2 Ambulance response time by ambulance zone (Non-Emergency)	67
RST-3 Transport of patients to hospital	69
PUB-1 Out-of-hospital cardiac arrests receiving bystander (non-EMS personnel) CPR ...	71

STATUTORY AUTHORITY

The California EMS Authority (EMSA or authority) is charged with creating a “statewide system for emergency medical services” and the responsibility for the “coordination and integration of all state activities concerning emergency medical services” (HS 1797.1). Moreover, the authority is required to assess each EMS area or the system’s service area, utilizing regional and local information, for “the purpose of determining the need for additional emergency medical services, coordination of emergency medical services and the effectiveness of emergency medical services” (HS1797.102). And local EMS agencies are required to plan, implement, and evaluate an EMS system (HS 1797.204).

Health and Safety Code 1797.103 identifies that one of the required elements of an EMS system is data collection and evaluation. Additionally, the development of quality improvement guidelines must be established (HS 1797.174). As a result of this statutory mandate, EMSA has developed regulations requiring the system data collection and evaluation, collection of prehospital care reports (CCR, Title 22, Division 9, Chapter 4, Section 100147, 100169, 100170).

Additionally, EMS system quality improvement regulations have been established (CCR, Title 22, Division 9, Chapter 12) that define the requirements for local EMS agencies, EMS service providers, and base hospitals in their role as part of the EMS system. These requirements include, but are not limited to the implementation of an EMSA approved EMS Quality Improvement program (EMS QI) and the use of defined indicators to assess the local EMS system as found in EMSA #166, Appendix E. This evaluation and EMS QI information must be submitted annually to EMSA, as part of its required EMS plan (HS 1797.254), in order to allow EMSA to evaluate if the plan effectively meets the needs of the persons served.

A report to the Legislature must be made on the effectiveness of EMS systems annually related to the EMS system’s impact on death and disability (HS 1797.121).

In order to achieve this mandate to evaluate system impact on patients, the continuum of care from dispatch to pre-hospital to hospital disposition must be connected. Only in this way, we can begin to understand how care provided by EMS personnel translates to improved outcomes and system effectiveness.

PROJECT HISTORY

The purpose of the EMS system core measures project is to increase the accessibility and accuracy of pre-hospital data for public, policy, academic and research purposes to facilitate EMS system evaluation and improvement through a grant from the California Health Care Foundation (CHCF). Ultimately, the project highlights opportunities to improve the quality of patient care delivered within an EMS system.

During a 1 year period, from April 1, 2012 to April 30, 2013, The California EMS Authority (EMSA) is performing the following activities to deliver a set of publicly available data reports:

1. Assess the capacity of the California Emergency Medical Services Information System to deliver core performance measures.
2. Create a formal data system profile and written analysis to identify areas for data quality improvement and inform an action plan to address the issues.
3. Work to reveal opportunities for both short-term and long-term data improvement plans.
4. Focus on achieving reliable measures that are high value and feasible within a short-term time frame.
5. Define and publish core measure sets that describe the coordination and effectiveness of EMS utilizing regional and local information for California. This project focuses upon the following core measure sets:
 - Trauma
 - Acute Coronary Syndrome/Heart Attack
 - Cardiac Arrest
 - Stroke
 - Respiratory
 - Pain Intervention
 - Pediatric
 - Skill Performance by EMS Providers
 - EMS Response and Transport
 - Public Education Bystander CPR
6. Submit California EMS data to the National EMS Information System (NEMSIS) to evaluate the statewide system from a national perspective.
7. Conduct three data workshops for local EMS agencies across the state to implement improved data collection and reporting practices with those Local Emergency Medical Services Agencies who participate in California Emergency Medical Services Information System.

WHAT ARE CORE MEASURES?

They are the use of standardized – or core – performance measures or quality indicators in examining an EMS system or treating an identified patient condition.

CORE MEASURES DEFINITION

The preliminary California EMS Core Measures were derived largely from a set of quality indicators developed through a project by the National Quality Forum. Additionally, NHSTA has published Performance Measures for emergency medical services. These California core measures will begin to benchmark the performance of EMS systems, perform recommended treatments determined to get the best results for patients with certain medical conditions, and transport patients to the most appropriate hospital. Information about these treatments are taken from the pre-hospital care reports and converted into a percentage.

The measures are based on scientific evidence about processes and treatments that are known to get the best results for a condition or illness. Core Measures help emergency medical services systems improve the quality of patient care by focusing on the actual results of care.

COMPARING PERFORMANCE

Emergency medical services systems across the state will be measured and compared on their performance in these Core Measures. There will be a delay between when data is reported from EMS systems and when it is available for review. This is because EMSA will have to wait for all local systems in the state to be compiled before it can post its quality data for a given period. This way, EMS systems and consumers can compare California program from the same time period.

In the future, EMS providers should utilize these core measures to assist in continuous quality improvement activities.

SYSTEM EVALUATION

The recurring theme in evaluation of the EMS system using these core measures consists of:

- Arrival at the scene in a timely manner,
- Timely, focused patient assessment,
- Delivery of time-sensitive prehospital therapy, and
- Transport to a hospital capable of providing necessary care

FUTURE CORE MEASURES

It is anticipated that the proposed EMS system cores measures may be modified and future core measures added in the future.

CORE MEASURES TASK FORCE

A task force has been convened to review the core measures and make recommendations. The task force consists of key data and quality leaders from local EMS agencies, medical directors, hospitals, and pre-hospital EMS providers.

QUALIFYING DATA

The data derived for all measures will come from the calendar years of 2009, 2010, and 2011. Reports will be run by calendar year to obtain longitudinal comparisons.

STANDARD ELEMENTS FOR EVERY MEASURE

The following standard elements are necessary to sort by time and location:

- Date/Time E05_01
- County E08_13
- LEMSA C01_01

REFERENCE INFORMATION

The California EMS System Core Quality Measures contains various references and coding from other documents. All data elements and values referenced in the Core Measures are coded using CEMSIS data standards, except where otherwise specified as NEMSIS. Please refer to the following documents regarding the codes found in each measure:

- EMSA #164: CEMSIS Data System Standards

NEMSIS 2.2.1 Data Dictionary – Updated 4/9/2012

NHTSA: Emergency Medical Services Performance Measures – Updated 12/2009
(www.ems.gov/pdf/811211.pdf)

EMS SYSTEM CORE MEASURES FOR CALIFORNIA

CCR Title 22, Div 9, Chap 12 100404	SET NAME	SET ID	PERFORMANCE MEASURE NAME	YEAR BEGIN TO BE MEASURED
D Clinical Care and Patient Outcome	Trauma (n=2)	TRA-1	Scene time for severely injured trauma patients	2013
		TRA-2	Direct transport to trauma center for severely injured trauma patients meeting criteria	2013
	Acute Coronary Syndrome (n=5)	ACS-1	Aspirin administration for chest pain/discomfort	2013
		ACS-2	12 lead EKG performance	2013
		ACS-3	Scene time for suspected heart attack patients	2013
		ACS-4	Advance hospital notification for suspected acute coronary syndrome	2014
		ACS-5	Direct transport to PCI center for suspected acute coronary syndrome (ACS) patients meeting criteria	2013
	Cardiac Arrest (n=4)	CAR-1	AED application prior to EMS Arrival	2014
		CAR-2	Out-of-hospital cardiac arrests return of spontaneous circulation	2013
		CAR-3	Out-of-hospital cardiac arrests survival to emergency department discharge	2013
		CAR-4	Out-of-hospital cardiac arrests survival to hospital discharge	2013
	Stroke (n=5)	STR-1	Identification of suspected stroke in the field	2014
		STR-2	Glucose testing for suspected stroke patients	2013
		STR-3	Scene time for suspected stroke patients	2013
		STR-4	Advance hospital notification for suspected stroke	2014
		STR-5	Direct transport to stroke center for suspected stroke patients meeting criteria	2013
	Respiratory (n=2)	RES-1	CPAP given for patients with respiratory distress	2014
		RES-2	Beta2 agonist administration	2013
	Pediatric (n=2)	PED-1	Pediatric asthma patients receiving bronchodilators	2013
		PED-2	Transport to pediatric trauma center	2014
	Pain Intervention (n=2)	PAI-1	Pain intervention	2013
		PAI-2	Results of pain intervention	2014

(Continued)

6 • California EMS System Core Quality Measures

CCR Title 22, Div 9, Chap 12 100404	SET NAME	SET ID	PERFORMANCE MEASURE NAME	YEAR BEGIN TO BE MEASURED
E Skills Maintenance and Competency	Performance of Skills (n=2)	SKL-1	Endotracheal intubation success rate	2013
		SKL-2	End-Tidal CO2 performed on any successful endotracheal intubation	2013
F Transportation and Facilities	Response and Transport (n=3)	RST-1	Ambulance response time by ambulance zone (Emergency)	2013
		RST-2	Ambulance response time by ambulance zone (Non-Emergency)	2013
		RST-3	Transport of patients to hospital	2013
G Public Education	Cardiopulmonary Resuscitation (n=1)	PUB-1	Out-of-hospital cardiac arrests receiving bystander (non-EMS Personnel/Responder) CPR	2014

Core Measures Specification Sheets

SCENE TIME FOR SEVERELY INJURED TRAUMA PATIENTS

MEASURE SET	Trauma	
SET MEASURE ID #	TRA-1	
PERFORMANCE MEASURE NAME	Scene time for severely injured trauma patients	
Description	On-Scene Time (90 th percentile) of severely injured Trauma Patients who were transported from the scene by ambulance	
Type of Measure	Process	
Reporting Value and Units	Time (Minutes and Seconds)	
Continuous Variable Statement (Population)	Time (in minutes) from time ambulance arrives at the scene until the time ambulance departs from the scene for Trauma patients, meeting criteria for transport to a trauma center (using revised trauma score or RTS<5), who received transport by ambulance to a hospital by EMS personnel (EMT, AEMT, and Paramedic).	
Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> • All events for which E02_04 “type of service requested” has value 30 “911 response (scene),”; and vehicle type corresponds to ground ambulance; • E02_20 “response mode to scene” has a value of 390 “lights and sirens” • Values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and logical; • Patients with E09_15 “provider primary impression” value 1740 “blunt injury” or 1741 “penetrating injury”, <u>or</u> E09_16 “provider secondary impression” value 1875 “blunt injury” or 1876 “penetrating injury” <u>and</u>: • patients with E14_27 “Revised Trauma Score” <5; <p><u>OR</u></p> <ul style="list-style-type: none"> • All events for which E02_04 “type of service requested” has value 30 “911 response (scene),”; and vehicle type corresponds to ground 	<ul style="list-style-type: none"> • Type of Service Requested (E02_04) • Response mode to scene (E02_20) • Arrived at Scene (E05_06) • Unit Left Scene (E05_09) • Provider Primary Impression (E09_15) • Provider Secondary Impression (E09_16) • Revised Trauma Score (E14_27) • Systolic Blood Pressure (E14_04) • Total GCS Value (E14_19) • Respiratory Rate (E14_11) • Date of Birth (E06_16) • Age Units (E06_15) • Age (E06_14)

	<p>ambulance; and E02_20 “response mode to scene” has a value of 390 “lights and sirens” and values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and logical;</p> <ul style="list-style-type: none"> • Patients with E09_15 “provider primary impression” values 1740 “blunt injury” or 1741 “penetrating injury”, or E09_16 “provider secondary impression” values 1875 “blunt injury” or 1876 “penetrating injury” <u>and:</u> • E14_19 “Total Glasgow Coma Score” value < 14; or • E14_04 “systolic blood pressure” value < 90; or • E14_11 “respiratory rate” value < 10 or > 29 for patients aged 1 year or older or E14_11 “respiratory rate” value < 20 for patients less than 1 year of age 	
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is the 90 th Percentile of the given numbers or distribution in their ascending order.	
Example of Final Reporting Value (number and units)	14 minutes, 34 seconds (14:34)	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format &	Process control or run chart by month	

10 • California EMS System Core Quality Measures

Frequency	
Suggested Statistical Measures	90 th Percentile Measurement. Aggregate measure of central tendency and quantile (fractile) measurement to determine the span of frequency distributions.
Trending Analysis	Yes
Benchmark Analysis	(TBD)
Rationale for Data	<input type="checkbox"/> The rapid transport of severe trauma patients from the scene to trauma centers has been well documented as a key indicator of survival.
References	NEMSIS Core Measure, Indicator 10.4

DIRECT TRANSPORT TO TRAUMA CENTER FOR SEVERELY INJURED TRAUMA PATIENTS MEETING CRITERIA

MEASURE SET	Trauma	
SET MEASURE ID #	TRA-2	
PERFORMANCE MEASURE NAME	Direct transport to trauma center for severely injured trauma patients meeting criteria	
Description	Trauma Patients who were transported from the scene directly to a Trauma Center	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	All trauma patients, meeting trauma criteria (using a Revised Trauma Score or RTS<5) for transport from scene to a trauma center	
Denominator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> • All events for which E02_04 “type of service requested” has value 30 “911 response (scene),”; and vehicle type corresponds to ground ambulance; • E02_20 “response mode to scene” has a value of 390 “lights and sirens” • Values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and logical; • Patients with E09_15 “provider primary impression” value 1740 “blunt injury” or 1741 “penetrating injury”, <u>or</u> E09_16 “provider secondary impression” value 1875 “blunt injury” or 1876 “penetrating injury” <u>and</u>: • patients with E14_27 “Revised Trauma Score” <5; <p><u>OR</u></p> <ul style="list-style-type: none"> • All events for which E02_04 “type of service requested” has value 30 “911 response (scene),”; and vehicle type corresponds to ground ambulance; and E02_20 “response mode to scene” has a value of 390 “lights and sirens” and values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and logical; 	<ul style="list-style-type: none"> • Provider Primary Impression (E09_15) • Provider Secondary Impression (E09_16) • Type of Service Requested (E02_04) • Revised Trauma Score (E14_27) • Systolic Blood Pressure (E14_04) • Total GCS Value (E14_19) • Respiratory Rate (E14_11) • Date of Birth (E06_16) • Age Units (E06_15) • Age (E06_14)

12 • California EMS System Core Quality Measures

	<ul style="list-style-type: none"> Patients with E09_15 “provider primary impression” values 1740 “blunt injury” or 1741 “penetrating injury”, or E09_16 “provider secondary impression” values 1875 “blunt injury” or 1876 “penetrating injury” <u>and</u>: E14_19 “Total Glasgow Coma Score” value < 14; or E14_04 “systolic blood pressure” value < 90; or E14_11 “respiratory rate” value < 10 or > 29 for patients aged 1 year or older or E14_11 “respiratory rate” value < 20 for patients less than 1 year of age 	
Exclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> All patients who were not transported to trauma center 	
Numerator Statement (sub-population)	Trauma patients, meeting criteria for transport to a trauma center, who received transport by ambulance directly to a trauma center by Ambulance	
Numerator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> All events for which E02_04 “type of service requested” has value 30 “911 response (scene),”; and vehicle type corresponds to ground ambulance; and E02_20 “response mode to scene” has a value of 390 “lights and sirens” and values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and logical; Patients with E09_15 “provider primary impression” value 1740 “blunt injury” or 1741 “penetrating injury”, or E09_16 “provider secondary impression” value 1875 “blunt injury” or 1876 “penetrating injury” <u>and</u>: patients with E14_27 “Revised Trauma Score” <5; <u>And</u> Patients who have “destination/transferred to” code (E20_02) of a trauma center <p><u>OR</u></p>	<ul style="list-style-type: none"> Revised Trauma Score (E14_27) Incident/Patient Disposition (E20_10) Hospital Destination (E20_02)

	<ul style="list-style-type: none"> • All events for which E02_04 “type of service requested” has value 30 “911 response (scene),”; and vehicle type corresponds to ground ambulance; and E02_20 “response mode to scene” has a value of 390 “lights and sirens” and values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and logical; • Patients with E09_15 “provider primary impression” values 1740 “blunt injury” or 1741 “penetrating injury”, or E09_16 “provider secondary impression” values 1875 “blunt injury” or 1876 “penetrating injury” <u>and:</u> • E14_19 “Total Glasgow Coma Score” value < 14; or • E14_04 “systolic blood pressure” value < 90; or • E14_11 “respiratory rate” value < 10 or > 29 for patients aged 1 year or older or E14_11 “respiratory rate” value < 20 for patients less than 1 year of age <u>And</u> • Patients who have “destination/transferred to” code (E20_02) of a trauma center 	
Exclusion Criteria	Criteria	Data Elements
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is N/D =%	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	

14 • California EMS System Core Quality Measures

Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.
Suggested Display Format & Frequency	Process control or run chart by month
Suggested Statistical Measures	Mean (x); Mode (m)
Trending Analysis	Yes
Benchmark Analysis	(TBD)
Rationale for Data	<input type="checkbox"/> The rapid transport of trauma patients to trauma centers has been well documented as a key indicator of survival.
References	NEMSIS Core Measure, Indicator 5

ASPIRIN ADMINISTRATION FOR CHEST PAIN/DISCOMFORT RATE

MEASURE SET	Acute Coronary Syndrome (ACS)	
SET MEASURE ID #	ACS-1	
PERFORMANCE MEASURE NAME	Aspirin administration for chest pain/discomfort rate	
Description	Patients aged 35 years and older with suspected cardiac chest pain who received aspirin by EMS personnel	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Improvement Noted As	An increase in the rate in terms of the percentage	
Denominator Statement (population)	Number of patients over age 35 creating a provider impression of chest pain/discomfort who are eligible for aspirin administration	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients with E09_15 1650 “Chest pain – suspected cardiac origin” or E09_16 value 1785 “chest pain – suspected cardiac origin”; Patients aged 35 years and older 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Age (E06_14) Age Units E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	Number of patients creating a provider impression of chest pain/discomfort who are eligible for and receive aspirin administration	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>

16 • California EMS System Core Quality Measures

	<ul style="list-style-type: none"> Patients with E09_15 1650 “Chest pain – suspected cardiac origin “or E09_16 value 1785 “chest pain – suspected cardiac origin”; Patients aged 35 years and older <u>And</u> E18_03 “medications given” equal to 8625 “aspirin” 	<ul style="list-style-type: none"> Medications given (E18_03)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is N/D =%	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	

<p>Rationale for Data</p>	<p>□ The early use of aspirin in patients with acute myocardial infarction results in a significant reduction in adverse events and subsequent mortality. The benefits of aspirin therapy on mortality are comparable to fibrinolytic therapy. The combination of aspirin and fibrinolytics provides additive benefits for patients with ST-elevation myocardial infarction (ISIS-2, 1988).</p> <p>□ Aspirin is also effective in patients with non-ST-elevation myocardial infarction (Theroux, 1988 and RISC Group, 1990). National guidelines strongly recommend early aspirin for patients hospitalized with AMI (Antman, 2004; Antman, 2008; and Wright, 2011).</p>
<p>References</p>	<p>NEMSIS Core Measure, Indicator 8</p>

12 LEAD ECG PERFORMANCE

MEASURE SET	Acute Coronary Syndrome (ACS)	
SET MEASURE ID #	ACS-2	
PERFORMANCE MEASURE NAME	12 Lead ECG Performance	
Description	Acute Coronary Syndrome (ACS) patients who received ECG by Paramedics	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	Number of patients creating a provider impression of chest pain/discomfort	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients with E09_15 1650 Chest pain – suspected cardiac origin or E09_16 value 1785 “chest pain – suspected cardiac origin”; Patients aged 35 years and older 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	Number of patients creating a provider impression of chest pain/discomfort who have 12-lead EKG performed	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients with E09_15 1650 Chest pain – suspected cardiac origin or E09_16 value 1785 “chest pain – suspected cardiac origin”; and Patients aged 35 years and older <u>and</u> Have a E19_03 “procedure” value 89.820 “12 lead -(Obtain)” or 89.821 “12 Lead (Transmitted) 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16) Procedures Performed (E19_03)

Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> Guidelines recommend patients presenting with chest discomfort or symptoms suggestive of ST-segment elevation myocardial infarction (STEMI) have a 12-lead electrocardiogram (ECG) performed within a target of 10 minutes of emergency department arrival (Krumholz, 2008). Evidence supports reperfusion benefits patients with identified STEMI (Antman 2004). <input type="checkbox"/> The diagnosis and management of STEMI patients is dependent upon practices within the emergency department. Timely ECGs assist in identifying STEMI patients and impact the choice of reperfusion strategy (Peacock, 2007). This measure will identify if 12-lead ECG for chest pain for AMI patients are being performed.	
References	NEMSIS Core Measure Indicator 7	

SCENE TIME FOR SUSPECTED HEART ATTACK PATIENTS

MEASURE SET	Acute Coronary Syndrome	
SET MEASURE ID #	ACS-3	
PERFORMANCE MEASURE NAME	Scene time for suspected heart attack patients	
Description	Scene Time of patients with field 12 lead ECG indicated ST Elevation Myocardial Infarction (STEMI)	
Type of Measure	Process	
Reporting Value and Units	Minutes	
Continuous Variable Statement (Population)	The 90 th percentile time interval in an emergency from EMS “arrived at scene” to “unit left scene”, for a given period of time, of patients having a recorded “STEMI” value for an indicator like E14_03 “cardiac rhythm”	
Denominator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> • All events for which E02_04 “type of service requested” has value 30 “911 response (scene),”; and vehicle type corresponds to ground ambulance; and • E02_20 “response mode to scene” has a value of 390 “lights and sirens” • Values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and logical; • Patients aged 35 years and older • Patient has a “STEMI” value recorded for an indicator like E14_03 “cardiac rhythm”, such as 3005, 3010, 3015 	<ul style="list-style-type: none"> • Type of Service Requested (E02_04) • Arrived at Scene (E05_06) • Unit Left Scene (E05_09) • Cardiac Rhythm (E14_03) • Age (E06_14) • Age Units (E06_15) • Date of Birth (E06_16)
Exclusion Criteria	Criteria	Data Elements
	None	
Indicator Formula Numeric Expression	The formula is the 90 th Percentile of the given numbers or distribution in their ascending order.	
Example of Final Reporting Value (number and units)	14 minutes, 20 seconds (14:20)	
Sampling	Yes	

Aggregation	Yes
Blinded	Yes
Minimum Data Values	30
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.
Suggested Display Format & Frequency	Process control or run chart by month
Suggested Statistical Measures	Mean (x); Mode (m)
Trending Analysis	Yes
Benchmark Analysis	(TBD)
Rationale for Data	
References	NEMSIS Core Measure Indicator 10.4

ADVANCE HOSPITAL NOTIFICATION FOR SUSPECTED ACS

MEASURE SET	Acute Coronary Syndrome	
SET MEASURE ID #	ACS-4 (To Be Measured in 2014)	
PERFORMANCE MEASURE NAME	Advance hospital notification for suspected Acute Coronary Syndrome	
Description	Advance hospital notification by EMS personnel to the receiving hospital via radio or telephone of patients identified in the prehospital setting as experiencing a STEMI	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	Number of patients having a recorded “STEMI” value for an indicator like E14_03 “cardiac rhythm”	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients aged 35 years and older Patients having E14_03 “cardiac rhythm” recorded with a “STEMI” value, such as 30005, 3010, 3015 	<ul style="list-style-type: none"> Age (E06_14) Age Units (E06_15) Date of Birth (E06_16) Cardiac Rhythm (E14_03)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	Number of patients having a recorded “STEMI” value for an indicator like E14_03 “cardiac rhythm” that have an E20_02 “destination/transferred to code” of an interventional cardiac cath center (STEMI Center) and advance hospital notification/activation/alert occurred	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients aged 35 years and older Patients having E14_03 “cardiac rhythm” recorded with a “STEMI” value, such as 30005, 3010, 3015 <u>And</u> have an E20_02 “destination/transferred to code” of (STEMI Center) an interventional cardiac cath center; and have hospital received advance 	<ul style="list-style-type: none"> Age (E06_14) Age Units (E06_15) Date of Birth (E06_16) Cardiac Rhythm (E14_03) Destination/Transferred to Code (E20_02) Hospital STEMI Alert Notification

	notification/activation/alert	
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> Advance hospital notification by EMS personnel may reduce the time to receive time sensitive diagnostics and therapy upon arrival at the emergency department. <input type="checkbox"/> Improved access to diagnostic imaging assists clinicians in the decision making process and treatment plans.	
References	CEMSIS Core Measures	

DIRECT TRANSPORT TO PCI CENTER FOR SUSPECTED ACUTE CORONARY SYNDROME (ACS) PATIENTS MEETING CRITERIA

MEASURE SET	Acute Coronary Syndrome	
SET MEASURE ID #	ACS-5	
PERFORMANCE MEASURE NAME	Direct transport to PCI center for suspected acute coronary syndrome (ACS) patients meeting criteria	
Description	Suspected Acute Coronary Syndrome (ACS) patients who were transported from the scene directly to a STEMI Center capable of PCI	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	Number of patients having a recorded “STEMI” value for an indicator like E14_03 “cardiac rhythm”	
Denominator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> Patients aged 35 years and older Patients having E14_03 “cardiac rhythm” recorded with a “STEMI” value, such as 30005, 3010, 3015 	<ul style="list-style-type: none"> Age (E06_14) Age Units (E06_15) Date of Birth (E06_16) Cardiac Rhythm (E14_03)
Exclusion Criteria		
	None	
Numerator Statement (sub-population)	Number of patients having a recorded “STEMI” value for an indicator like E14_03 “cardiac rhythm” that have an E20_02 “destination/ transferred to code” of an interventional cardiac cath center (STEMI Center)	
Numerator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> Patients aged 35 years and older Patients having E14_03 “cardiac rhythm” recorded with a “STEMI” value, such as 30005, 3010, 3015 <u>And</u> that have an E20_02 “destination/transferred to 	<ul style="list-style-type: none"> Age (E06_14) Age Units (E06_15) Date of Birth (E06_16) Cardiac Rhythm (E14_03) Cardiac Rhythm (E14_03) Destination/Transferred to Code (E20_02)

	code” of an interventional cardiac cath center (STEMI Center)	
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data		
References	NEMSIS Core Measure Indicator 9	

AED PRIOR TO EMS ARRIVAL

MEASURE SET	Cardiac Arrest	
SET MEASURE ID #	CAR-1 (To Be Measured in 2014)	
PERFORMANCE MEASURE NAME	AED prior to EMS arrival	
Description	Application by bystander (non-EMS personnel/responders) of AED to cardiac arrest patient prior to EMS Arrival	
Type of Measure	Process	
Reporting Value and Units	Numeric Value	
Denominator Statement (population)	All patients in Cardiac Arrest	
Denominator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> Patients having a recorded E11_01 “cardiac arrest” value of 2240 “yes, Prior to EMS arrival” or value of 2245 “yes, after EMS arrival”; E11_02 “cardiac arrest etiology” value of 2250 “presumed cardiac” E11_03 “resuscitation attempted” values 2280 “attempted defibrillation” or 2285 “attempted ventilation” or 2290 “initiated chest compressions” 	<ul style="list-style-type: none"> Cardiac Arrest (E11_01) Cardiac Arrest Etiology (E11_02) Resuscitation Attempted (E11_03)
Exclusion Criteria	Criteria	Data Elements
	None	
Numerator Statement (sub-population)	Patients in Cardiac Arrest who had AED application prior to arrival of EMS personnel	
Numerator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> Patients having a recorded E11_01 “cardiac arrest” value of 2240 “yes, Prior to EMS arrival” or value of 2245 “yes, after EMS arrival”; E11_02 “cardiac arrest etiology” value of 2250 “presumed cardiac” E11_03 “resuscitation attempted” values 2280 “attempted defibrillation” or 2285 “attempted 	<ul style="list-style-type: none"> Cardiac Arrest (E11_01) Cardiac Arrest Etiology (E11_02) Resuscitation Attempted (E11_03) Prior Aid Performed by (E09_02) Automated External Defibrillation (AED) Performed

	<p>ventilation” or 2290 “initiated chest compressions”</p> <p><u>And</u></p> <ul style="list-style-type: none"> • E09_02 “prior aid performed by” has a value of 1205 “lay person”, 1200 “law enforcement”, 1210 “other health care provider”, and • Automated External Defibrillation (AED) was performed. 	
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data		
References	CEMSIS Core Measures	

OUT-OF-HOSPITAL CARDIAC ARRESTS RETURN OF SPONTANEOUS CIRCULATION

MEASURE SET	Cardiac Arrest	
SET MEASURE ID #	CAR-2	
PERFORMANCE MEASURE NAME	Out-of-hospital cardiac arrests return of spontaneous circulation	
Description	Number of patients experiencing cardiac origin cardiac arrest who have Return of Spontaneous Circulation (ROSC) at any time (Utstein) in a given period	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	Total number of patients in a given period experiencing cardiac origin cardiac arrest	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients having a recorded E11_01 “cardiac arrest” value of 2240 “yes, Prior to EMS arrival” or value of 2245 “yes, after EMS arrival”; E11_02 “cardiac arrest etiology” value of 2250 “presumed cardiac” E11_03 “resuscitation attempted” values 2280 “attempted defibrillation” or 2285 “attempted ventilation” or 2290 “initiated chest compressions” 	<ul style="list-style-type: none"> Cardiac Arrest (E11_01) Cardiac Arrest Etiology (E11_02) Resuscitation Attempted (E11_03) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	Number of patients experiencing cardiac origin cardiac arrest who have a return of spontaneous circulation (ROSC)	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients having a recorded E11_01 “cardiac arrest” value of 2240 “yes, Prior to EMS arrival” or value of 2245 “yes, after EMS arrival”; 	<ul style="list-style-type: none"> Cardiac Arrest (E11_01) Cardiac Arrest Etiology (E11_02) Resuscitation Attempted (E11_03)

	<ul style="list-style-type: none"> E11_02 “cardiac arrest etiology” value of 2250 “presumed cardiac” E11_03 “resuscitation attempted” values 2280 “attempted defibrillation” or 2285 “attempted ventilation” or 2290 “initiated chest compressions” <u>And</u> E11_06 “any return of spontaneous circulation” values 2370 “yes, prior to ED Arrival Only” or 2375 “yes, prior to ED arrival and at the ED” 	<ul style="list-style-type: none"> Any Return to Spontaneous Circulation (E11_06)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is N/D =%	
Example of Final Reporting Value (number and units)	25%	
Sampling	No	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> Definitive care for ACS	
References	NEMSIS Core Measure Indicator 18	

OUT-OF-HOSPITAL CARDIAC ARRESTS SURVIVAL TO ED DISCHARGE

MEASURE SET	Cardiac Arrest	
SET MEASURE ID #	CAR-3	
PERFORMANCE MEASURE NAME	Out-of-hospital Cardiac Arrests Survival to ED discharge	
Description	Number of patients experiencing cardiac origin cardiac arrest after EMS arrival who survive to discharge from the ED divided by the total number of patients experiencing cardiac origin cardiac arrest in a given period	
Type of Measure	Outcome	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	Total number of patients experiencing cardiac origin cardiac arrest in a given period	
Denominator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> Patients having a recorded E11_01 “cardiac arrest” value of 2240 “yes, Prior to EMS arrival” or value of 2245 “yes, after EMS arrival”; E11_02 “cardiac arrest etiology” value of 2250 “presumed cardiac” E11_03 “resuscitation attempted” values 2280 “attempted defibrillation” or 2285 “attempted ventilation” or 2290 “initiated chest compressions” 	<ul style="list-style-type: none"> Cardiac Arrest (E11_01) Cardiac Arrest Etiology (E11_02) Resuscitation Attempted (E11_03) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	Criteria	Data Elements
	None	
Numerator Statement (sub-population)	Number of patients experiencing cardiac origin cardiac arrest who survive to discharge from the ED	
Numerator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> Patients having a recorded E11_01 “cardiac arrest” value of 2240 “yes, Prior to EMS arrival” or value of 2245 “yes, after EMS arrival”; E11_02 “cardiac arrest etiology” 	<ul style="list-style-type: none"> Cardiac Arrest (E11_01) Cardiac Arrest Etiology (E11_02) Resuscitation Attempted (E11_03) Emergency Department

	<p>value of 2250 “presumed cardiac”</p> <ul style="list-style-type: none"> E11_03 “resuscitation attempted” values 2280 “attempted defibrillation” or 2285 “attempted ventilation” or 2290 “initiated chest compressions” <p><u>And</u></p> <ul style="list-style-type: none"> E22_01 “emergency department disposition” values 5335 “admitted to hospital floor” or 5340 “admitted to hospital ICU” or 5355 “released” or 5360 “transferred” 	Disposition (E22_01)
Exclusion Criteria	Criteria	Data Elements
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	25%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> Cardiac Arrest survival has been shown to increase with early CPR and Automated External Defibrillation	
References	NEMIS Core Measure Indicator 18.1	

OUT-OF-HOSPITAL CARDIAC ARRESTS SURVIVAL TO HOSPITAL DISCHARGE

MEASURE SET	Cardiac Arrest	
SET MEASURE ID #	CAR-4	
PERFORMANCE MEASURE NAME	Out-of-hospital Cardiac Arrests Survival to hospital discharge	
Description	Number of patients experiencing cardiac origin cardiac arrest after EMS arrival who survive to discharge from the hospital divided by the total number of patients experiencing cardiac origin cardiac arrest	
Type of Measure	Outcome	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	Total number of patients experiencing cardiac origin cardiac arrest in a given period	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients having a recorded E11_01 “cardiac arrest” value of 2240 “yes, Prior to EMS arrival” or value of 2245 “yes, after EMS arrival”; E11_02 “cardiac arrest etiology” value of 2250 “presumed cardiac” E11_03 “resuscitation attempted” values 2280 “attempted defibrillation” or 2285 “attempted ventilation” or 2290 “initiated chest compressions” 	<ul style="list-style-type: none"> Cardiac Arrest (E11_01) Cardiac Arrest Etiology (E11_02) Resuscitation Attempted (E11_03) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	Number of patients experiencing cardiac origin cardiac arrest who survive to discharge from the hospital	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients having a recorded E11_01 “cardiac arrest” value of 2240 “yes, Prior to EMS arrival” or value of 2245 “yes, after EMS arrival”; E11_02 “cardiac arrest etiology” value of 2250 “presumed 	<ul style="list-style-type: none"> Cardiac Arrest (E11_01) Cardiac Arrest Etiology (E11_02) Resuscitation Attempted (E11_03) Hospital Disposition

	cardiac” <ul style="list-style-type: none"> E11_03 “resuscitation attempted” values 2280 “attempted defibrillation” or 2285 “attempted ventilation” or 2290 “initiated chest compressions” <u>And</u> <ul style="list-style-type: none"> E22_02 “hospital disposition” values 5370 “discharged” or 5375 “transfer to hospital” or 5380 “transfer to nursing home” or 5385 “transfer to other” or 5390 “transfer to rehabilitation facility” 	(E22_02)
Exclusion Criteria	Criteria	Data Elements
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	25%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> Cardiac Arrest survival has been shown to increase with early CPR and Automated External Defibrillation	
References	NEMSIS Core Measure Indicator 18.2	

IDENTIFICATION OF SUSPECTED STROKE IN THE FIELD

MEASURE SET	Stroke	
SET MEASURE ID #	STR-1 (To Be Measured in 2014)	
PERFORMANCE MEASURE NAME	Identification of Suspected Stroke in the Field	
Description	Identification of suspected stroke by EMS personnel in the field using a validated pre-hospital stroke screen (Los Angeles LAPSS or Cincinnati CPSS)	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	Patients with a Provider Primary Impression of Suspected Stroke	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> • Patients with E09_15 value 1730 value “neurological deficit (includes CVA/TIA)” or E09_16 value 1865 “neurological deficit (includes CVA/TIA)” • Patients aged 18 years of age or older 	<ul style="list-style-type: none"> • Provider Primary Impression (E09_15) • Provider Secondary Impression (E09_16) • Age (E06_14) • Age Units (E06_15) • Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	Identification of a suspected stroke used a validated pre-hospital stroke screen (Los Angeles LAPSS or Cincinnati CPSS)	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> • Patients with E09_15 value 1730 value “neurological deficit (includes CVA/TIA)” or E09_16 value 1865 “neurological deficit (includes CVA/TIA)” • Patients aged 18 years of age or older <li style="padding-left: 20px;"><u>And</u> • Used a validated pre-hospital stroke screen, such as indicated by NEMSIS E16_24 “Neurological Assessment” values if available 	<ul style="list-style-type: none"> • Provider Primary Impression (E09_15) • Provider Secondary Impression (E09_16) • Validated Pre-Hospital Stroke Screen (E16_24)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>

	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> Prehospital assessment and advance hospital notification by EMS personnel may reduce the time to receive time sensitive diagnostics and therapy upon arrival at the emergency department. <input type="checkbox"/> Improved access to diagnostic imaging assists clinicians in the decision making process and treatment plans.	
References	CEMSIS Core Measure	

GLUCOSE TESTING FOR SUSPECTED STROKE PATIENTS

MEASURE SET	Stroke	
SET MEASURE ID #	STR-2	
PERFORMANCE MEASURE NAME	Glucose Testing for Suspected Stroke patients	
Description	Patients with suspected stroke have assessment of blood glucose level.	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	All Suspected Stroke patients	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients with E09_15 value 1730 value “neurological deficit (includes CVA/TIA)” or E09_16 value 1865 “neurological deficit (includes CVA/TIA)” Patients aged 18 years of age or older 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	Glucose level Checked on all suspected Stroke patients	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients with E09_15 value 1730 value “neurological deficit (includes CVA/TIA)” or E09_16 value 1865 “neurological deficit (includes CVA/TIA)” Patients aged 18 years of age or older <u>And</u> Patient received glucose testing E19_03 “procedure” with a value of 38.995 “blood glucose analysis” 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Procedure (E19_03)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>

	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> <input type="checkbox"/>	
References	CEMSIS Core Measures	

SCENE TIME FOR SUSPECTED STROKE PATIENTS

MEASURE SET	Stroke	
SET MEASURE ID #	STR-3	
PERFORMANCE MEASURE NAME	Scene time for suspected stroke patients	
Description	Scene Time for Suspected Stroke Patients who were transported from the scene by ambulance	
Type of Measure	Process	
Reporting Value and Units	Minutes	
Continuous Variable Statement (population)	All suspected stroke patients	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> All events for which E02_04 “type of service requested” has value 30 “911 response (scene),” vehicle type corresponds to ground ambulance; and Values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and pass logic test; Patients with E09_15 value 1730 value “neurological deficit (includes CVA/TIA)” or E09_16 value 1865 “neurological deficit (includes CVA/TIA)” Patients aged 18 years of age or older 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Type of Service Requested (E02_04) Unit Arrived at Scene (E05_06) Unit Left Scene (E05_09) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is the 90 th Percentile of the given numbers or distribution in their ascending order.	
Example of Final Reporting Value (number and units)	14 minutes, 20 seconds (14:20)	
Sampling	Yes	
Aggregation	Yes	

Blinded	Yes
Minimum Data Values	30
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.
Suggested Display Format & Frequency	Process control or run chart by month
Suggested Statistical Measures	Mean (x); Mode (m)
Trending Analysis	Yes
Benchmark Analysis	(TBD)
Rationale for Data	<input type="checkbox"/> The rapid transport of suspected stroke patients to stroke centers which have CT or MRI scan capabilities to assist clinicians in the decision making process and treatment plan has been documented as a key indicator of the quality of care a patient receives.
References	CEMSIS Core Measures

ADVANCE HOSPITAL NOTIFICATION FOR SUSPECTED STROKE

MEASURE SET	Stroke	
SET MEASURE ID #	STR-4 (To Be Measured in 2014)	
PERFORMANCE MEASURE NAME	Advance hospital notification for suspected stroke	
Description	Advance hospital notification by EMS personnel to the receiving hospital via radio or telephone of patients identified in the prehospital setting as experiencing a suspected stroke, preferably using a validated pre-hospital stroke screen	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	Patients with a Provider Primary or Secondary Impression of Suspected Stroke	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients with E09_15 value 1730 value “neurological deficit (includes CVA/TIA)” or E09_16 value 1865 “neurological deficit (includes CVA/TIA)” Patients aged 18 years of age or older 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	Receiving Hospitals who received advance notification/alert by EMS personnel (EMT, AEMT, and Paramedic) for patients with Suspected Stroke patients	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients with E09_15 value 1730 value “neurological deficit (includes CVA/TIA)” or E09_16 value 1865 “neurological deficit (includes CVA/TIA)” Patients aged 18 years of age or older <u>And</u> Advance hospital Stroke notification/alert 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Hospital Disposition (E22_02) Advance Hospital Notification/Alert

Exclusion Criteria	Criteria	Data Elements
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> Improved access to diagnostic imaging assists clinicians in the decision making process and treatment plans. Over 143,579 people die each year from stroke (Stroke Center, 2009). Stroke is the third leading cause of death in the United States. Each year, about 795,000 people suffer a stroke. About 600,000 of these are first attacks, and 185,000 are recurrent attacks (AHA, 2009). Decreasing radiology turnaround times will enhance decision making capabilities for patients with TIA or Acute Ischemic Stroke. Of all strokes, 87 percent are ischemic, 10 percent are intracerebral hemorrhage, and 3 percent are subarachnoid hemorrhage (NINDS, 2004). Because of the therapeutic time window for treatment possibilities, timely completion and results of the CT or MRI scan are imperative and will directly impact the quality of care a patient receives. <input type="checkbox"/> Advance hospital notification by EMS personnel may reduce the time to receive time sensitive diagnostics and therapy upon arrival at the emergency department.	
References	CEMSIS Core Measures	

DIRECT TRANSPORT TO STROKE CENTER FOR SUSPECTED STROKE PATIENTS MEETING CRITERIA

MEASURE SET	Stroke	
SET MEASURE ID #	STR-5	
PERFORMANCE MEASURE NAME	Direct transport to stroke center for suspected stroke patients meeting criteria	
Description	Suspected Stroke Patients who were transported from the scene directly to a Stroke Center	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	All stroke patients, meeting local stroke criteria for transport to a stroke center	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients with E09_15 value 1730 value “neurological deficit (includes CVA/TIA)” or E09_16 value 1865 “neurological deficit (includes CVA/TIA)” Patients aged 18 years of age or older 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	Suspected Stroke patients, meeting criteria for transport to a stroke center, who received transport by ambulance directly to a stroke center	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients with E09_15 value 1730 value “neurological deficit (includes CVA/TIA)” or E09_16 value 1865 “neurological deficit (includes CVA/TIA)” Patients aged 18 years of age or older <u>And</u> E20_01 “Destination Transferred To, 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Destination/Transferred To (E20_01)

	Name" represents a stroke center	
Exclusion Criteria	Criteria	Data Elements
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> The rapid transport of suspected stroke patients directly to stroke centers has been well documented to improve access to diagnostic imaging and treatment. Prehospital assessment and advance hospital notification by EMS personnel also reduces the time to receive time sensitive diagnostics and therapy upon arrival at the emergency department. <input type="checkbox"/> Improved access to diagnostic imaging assists clinicians in the decision making process and treatment plans. Over 143,579 people die each year from stroke (Stroke Center, 2009). Stroke is the third leading cause of death in the United States. Each year, about 795,000 people suffer a stroke. About 600,000 of these are first attacks, and 185,000 are recurrent attacks (AHA, 2009). Of all strokes, 87 percent are ischemic, 10 percent are intracerebral hemorrhage, and 3 percent are subarachnoid hemorrhage (NINDS, 2004). Because of the therapeutic time window for treatment possibilities, timely completion and results of the CT or MRI scan are imperative and will directly impact the quality of care a patient receives.	
References	CEMSIS Core Measures	

CPAP GIVEN FOR PATIENTS WITH RESPIRATORY DISTRESS

MEASURE SET	Respiratory	
SET MEASURE ID #	RES-1 (To Be Measured in 2014)	
PERFORMANCE MEASURE NAME	CPAP given for patients with respiratory distress	
Description	Patients with respiratory distress that have CPAP used in their treatment.	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	Number of patients creating a provider impression of respiratory distress	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients for whom E09_15 “provider’s primary impression” has value 1701 “shortness of breath – suspected asthma/COPD” or 1702 “shortness of breath – suspected pulmonary edema” or for whom E09_16 “provider’s secondary impression” has value 1835 – “shortness of breath – suspected asthma/COPD” or 1836 “shortness of breath – suspected pulmonary edema” Patients aged 14 years or older 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	Number of patients creating a provider impression of respiratory arrest/distress who receive CPAP	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients for whom E09_15 “provider’s primary impression” has value 1701 “shortness of breath – suspected asthma/COPD” or 1702 “shortness of breath – suspected pulmonary edema” 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Procedure (NEMSIS E19_03)

	<p>or for whom E09_16 “provider’s secondary impression” has value 1835 – “shortness of breath – suspected asthma/COPD” or 1836 “shortness of breath – suspected pulmonary edema”</p> <ul style="list-style-type: none"> • Patients aged 14 years or older • <u>And</u> • who have a NEMESIS E19_03 value for 93.900 “CPAP” 	
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> <input type="checkbox"/>	
References	CEMSIS Core Measures	

BETA2 AGONIST ADMINISTRATION

Measure Set	Respiratory	
Set Measure ID #	RES-2	
Performance Measure Name	Beta2 agonist administration	
Description	Frequency of beta2 agonist (i.e., Albuterol) administered by EMS personnel for patients with signs and symptoms of shortness of breath and/or suspected bronchospasm.	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	Number of adult patients creating a provider impression of respiratory distress	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients for whom E09_15 “provider’s primary impression” has value 1701 “shortness of breath – suspected asthma/COPD” or 1702 “shortness of breath – suspected pulmonary edema” or for whom E09_16 “provider’s secondary impression” has value 1835 – “shortness of breath – suspected asthma/COPD” or 1836 “shortness of breath – suspected pulmonary edema” Patients aged 14 years or older 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	Adult patients with signs and symptoms of shortness of breath and bronchospasm who receive a beta agonist by EMS personnel in the prehospital setting.	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients for whom E09_15 “provider’s primary impression” has value 1701 “shortness of 	<ul style="list-style-type: none"> Provider Primary Impression (NEMESIS E09_15) Provider Secondary

	<p>breath – suspected asthma/COPD” or 1702 “shortness of breath – suspected pulmonary edema” or for whom E09_16 “provider’s secondary impression” has value 1835 – “shortness of breath – suspected asthma/COPD” or 1836 “shortness of breath – suspected pulmonary edema”</p> <ul style="list-style-type: none"> • Patients aged 14 years or older <u>And</u> • Who have a CEMSIS E18_03 value 8620 “aerosolized or nebulized beta-2 specific bronchodilator”, 8635 “Beta agonist”, or 8700 “Ipratropium Bromide”.; or a NEMSIS E18_03 element indicating any of the above 	<p>Impression (NEMSIS E09_16)</p> <ul style="list-style-type: none"> • Age (E06_14) • Age Units (E06_15) • Date of Birth (E06_16) • Medication Given (CEMSIS E18_03) • Medication Given (NEMSIS E18_03) • Medication Route (E18_04)
Exclusion Criteria	Criteria	Data Elements
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	

48 • California EMS System Core Quality Measures

Trending Analysis	Yes
Benchmark Analysis	(TBD)
Rationale for Data	<p><input type="checkbox"/> Short-acting selective beta agonists (e.g., albuterol, levalbuterol) are the first line of therapy for the quick relief of acute asthma symptoms (e.g., shortness of breath, cough, wheeze, chest tightness) Beta agonists medications help to reduce the symptoms of acute bronchospasm by relaxing the bronchial smooth muscle walls. http://www.uptodate (2012).</p> <p><input type="checkbox"/> Guidelines for the management of Chronic Obstructive Lung Disease (COPD) have been published by the Global Initiative for Chronic Obstructive Lung Disease (GOLD) and the American Thoracic Society-European Respiratory Society. Bronchodilator medications are an important pharmacologic intervention for persons with COPD. Prevention strategies have been developed for acute exacerbations, and management strategies include using short-acting beta2-agonists, systemic corticosteroids, and antibiotics, in both at home and hospital settings. Review of guidelines and the literature in the treatment of acute bronchospasm in chronic obstructive pulmonary disease. Pharmacotherapy. 2006 Sep;26(9 Pt 2):156S-63S</p>
References	CEMSIS Core Measures

PEDIATRIC ASTHMA PATIENTS RECEIVING BRONCHODILATORS

MEASURE SET	Pediatric	
SET MEASURE ID #	PED-1	
PERFORMANCE MEASURE NAME	Pediatric asthma patients receiving bronchodilators	
Description	Frequency of bronchodilators or beta2 agonists (i.e., Albuterol) administered by EMS personnel for patients younger than age 14 years with signs and symptoms of shortness of breath and/or suspected bronchospasm	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	All pediatric patients with respiratory distress from Asthma	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients for whom E09_15 “provider’s primary impression” has value 1701 “shortness of breath – suspected asthma/COPD” or for whom E09_16 “provider’s secondary impression” has value 1835 – “shortness of breath – suspected asthma/COPD” Patients less than 14 years of age 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	All pediatric patients with respiratory distress from Asthma receiving bronchodilators	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients for whom E09_15 “provider’s primary impression” has value 1701 “shortness of breath – suspected asthma/COPD” or for whom E09_16 “provider’s secondary impression” has value 1835 – 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)

50 • California EMS System Core Quality Measures

	<p>“shortness of breath – suspected asthma/COPD”</p> <ul style="list-style-type: none"> • Patients less than 14 years of age <p><u>And</u></p> <ul style="list-style-type: none"> • Who have a CEMSIS E18_03 value 8620 “aerosolized or nebulized beta-2 specific bronchodilator”, 8635 “Beta agonist”, or 8700 “Ipratropium Bromide”; or a NEMSIS E18_03 element indicating any of the above 	<ul style="list-style-type: none"> • Medication Given (CEMSIS E18_03) • Medication Given (NEMSIS E18_03)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is N/D =%	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> <input type="checkbox"/>	

References	<ol style="list-style-type: none">1. NEMSIS Core Measures2. CEMSIS Core Measures
-------------------	---

TRANSPORT TO PEDIATRIC TRAUMA CENTER

MEASURE SET	Pediatric	
SET MEASURE ID #	PED-2 (to be measured in 2014)	
PERFORMANCE MEASURE NAME	Transport to pediatric trauma center	
Description	Direct transport to pediatric trauma center for those severely injured pediatric patients, less than 14 years of age, meeting criteria	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	All pediatric Trauma patients	
Denominator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> All events for which E02_04 “type of service requested” has value 30 “911 response (scene),”; and vehicle type corresponds to ground ambulance; E02_20 “response mode to scene” has a value of 390 “lights and sirens” Values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and logical; Patients with E09_15 “provider primary impression” value 1740 “blunt injury” or 1741 “penetrating injury”, <u>or</u> E09_16 “provider secondary impression” value 1875 “blunt injury” or 1876 “penetrating injury” Patients less than 14 years old <u>and</u>: patients with E14_27 “Revised Trauma Score” <5; <p><u>OR</u></p> <ul style="list-style-type: none"> All events for which E02_04 “type of service requested” has value 30 “911 response (scene),”; and vehicle type corresponds to ground ambulance; and E02_20 	<ul style="list-style-type: none"> Provider Primary Impression (E09_15) Provider Secondary Impression (E09_16) Type of Service Requested (E02_04) Response mode to scene (E02_20) Arrived at Scene (E05_06) Unit left scene (E05_09) Revised Trauma Score (E14_27) Systolic Blood Pressure (E14_04) Total GCS Value (E14_19) Respiratory Rate (E14_11) Date of Birth (E06_16) Age Units (E06_15) Age (E06_14)

	<p>“response mode to scene” has a value of 390 “lights and sirens” and values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and logical;</p> <ul style="list-style-type: none"> • Patients with E09_15 “provider primary impression” values 1740 “blunt injury” or 1741 “penetrating injury”, or E09_16 “provider secondary impression” values 1875 “blunt injury” or 1876 “penetrating injury” • Patients less than 14 years old <u>and</u>: • E14_19 “Total Glasgow Coma Score” value < 14; or • E14_04 “systolic blood pressure” value < 90; or • E14_11 “respiratory rate” value < 10 or > 29 for patients aged 1 year or older or E14_11 “respiratory rate” value < 20 for patients less than 1 year of age 	
Exclusion Criteria	Criteria	Data Elements
	None	
Numerator Statement (sub-population)	Pediatric trauma patients transported to a pediatric trauma center	
Numerator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> • All events for which E02_04 “type of service requested” has value 30 “911 response (scene),”; and vehicle type corresponds to ground ambulance; • E02_20 “response mode to scene” has a value of 390 “lights and sirens” • Values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and logical; • Patients with E09_15 “provider primary impression” value 1740 “blunt injury” or 1741 “penetrating injury”, or E09_16 “provider secondary impression” 	<ul style="list-style-type: none"> • Provider Primary Impression (E09_15) • Provider Secondary Impression (E09_16) • Type of Service Requested (E02_04) • Response mode to scene (E02_20) • Arrived at Scene (E05_06) • Unit left scene (E05_09) • Revised Trauma Score (E14_27) • Systolic Blood Pressure (E14_04) • Total GCS Value (E14_19) • Respiratory Rate (E14_11)

	<p>value 1875 “blunt injury” or 1876 “penetrating injury”</p> <ul style="list-style-type: none"> • Patients less than 14 years old <u>and</u>: • patients with E14_27 “Revised Trauma Score” <5; And • Patients who have “destination/transferred to” code (E20_02) of a pediatric trauma center <p><u>OR</u></p> <ul style="list-style-type: none"> • All events for which E02_04 “type of service requested” has value 30 “911 response (scene),”; and vehicle type corresponds to ground ambulance; and E02_20 “response mode to scene” has a value of 390 “lights and sirens” and values for “arrived at scene” E05_06 and “unit left scene” E05_09 are present and logical; • Patients with E09_15 “provider primary impression” values 1740 “blunt injury” or 1741 “penetrating injury”, or E09_16 “provider secondary impression” values 1875 “blunt injury” or 1876 “penetrating injury” • Patients less than 14 years old <u>and</u>: • E14_19 “Total Glasgow Coma Score” value < 14; or • E14_04 “systolic blood pressure” value < 90; or • E14_11 “respiratory rate” value < 10 or > 29 for patients aged 1 year or older or E14_11 “respiratory rate” value < 20 for patients less than 1 year of age <u>And</u> • Patients who have “destination/transferred to” code (E20_02) of a pediatric trauma center 	<ul style="list-style-type: none"> • Date of Birth (E06_16) • Age Units (E06_15) • Age (E06_14) • Destination Transferred to (E20_02)
--	--	---

Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> <input type="checkbox"/>	
References	CEMSIS Core Measures	

PAIN INTERVENTION

MEASURE SET	Pain Intervention	
SET MEASURE ID #	PAI-1	
PERFORMANCE MEASURE NAME	Pain intervention	
Description	Percentage of patients age 14 and older reporting a pain value of 7 or greater on a 0-10 scale that received subsequent interventions associated with pain relief	
Type of Measure	Process	
Reporting Value and Units	Percentage	
Denominator Statement (Population)	The total number of events over a given period in which patients reported as having a pain value of 7 or greater during the continuum of the EMS call	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Events in which patients had recorded a pain value of 7 or greater for E14_23 Patient aged 14 years or older (E06_14) 	<ul style="list-style-type: none"> Pain Scale (E14_23) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients with no value recorded for NEMSIS E14_01, who have no value for either NEMSIS E18_01 or NEMSIS E19_01, to indicate the intervention occurred after pain measurement; 	<ul style="list-style-type: none"> Date Time Vitals Taken (NEMSIS E14_01) Date Time Medication Administered (NEMSIS E18_01) Date Time Procedure Performed Successfully (NEMSIS E19_01)
Numerator Statement (sub-population)	The total number of events over a given period in which patients reported as having a pain value of 7 or greater during the continuum of the EMS call	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Events in which patients had recorded a pain value of 7 or greater for E14_23 Patient aged 14 years or older (E06_14) <u>And</u> Associated value for NEMSIS E14_01, Who have at least one value for E18_03 or E19_03 representing 	<ul style="list-style-type: none"> Pain Scale (E14_23) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16) Date Time Vitals Taken (NEMSIS E14_01) Date Time Medication Administered (NEMSIS E18_01)

	a accepted intervention recognized for pain relief, and the related NEMSIS E18_01 or NEMSIS E19_01 elements indicate the interventions occurred after the pain scale was assessed.	<ul style="list-style-type: none"> • Medication Given (E18_03) • Procedure (E19_03)
Exclusion Criteria	Criteria	Data Elements
	Patients with no value recorded for NEMSIS E14_01 associated with administration of the pain scale E14_23; or who have no logical values for E18_01 or E19_01 to indicate the intervention occurred after assessment of pain scale ≥ 7	<ul style="list-style-type: none"> • Date Time Vitals Taken (NEMSIS E14_01) • Date Time Medication Administered (NEMSIS E18_01) • Date Time Procedure Performed Successfully (NEMSIS E19_01)
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> Administration of pain treatment is important to determine the effectiveness of medicine given to patient to reduce patient pain scale level	
References	NEMSIS Core Measure, Indicator 6.4	

RESULTS OF PAIN INTERVENTION

Measure Set	Pain Intervention	
Set Measure ID #	PAI-2 (to be measured in 2014)	
Performance Measure Name	Results of pain intervention	
Description	Percentage of patients age 14 and older reported decreased pain subsequent to interventions associated with pain relief when comparing first and last pain scale (0-10) values	
Type of Measure	Outcome	
Reporting Value and Units	Percentage	
Denominator Statement (Population)	The total number of events over a given period in which patients had at least two “measurements” of pain during the continuum of the EMS call	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Events in which patients had recorded at least two values for E14_23 each with a different associated value for NEMSIS E14_01 Patient aged 14 years or older (E06_14) Received Interventions for Pain Relief 	<ul style="list-style-type: none"> Pain Scale (E14_23) Age (E06_14) Age Units (E06_15) Date of Birth (E06_16) Date/Time Vital Taken (NEMSIS E14_01) Date Time Medication Administered (NEMSIS E18_01) Date Time Procedure Performed Successfully (NEMSIS E19_01)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Patients with one or no value recorded for NEMSIS E14_01, or who have at least two values for NEMSIS E14_23 but those values have no clear associated values for E14_01 or fail a logic test; 	
Numerator Statement (sub-population)	Number of events for a given period in which patients’ pain scale values decreased over the continuum of the EMS call	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>

	<ul style="list-style-type: none"> • Events in which patients had recorded at least two values for E14_23 each with a different associated value for NEMSIS E14_01 • Patient aged 14 years or older (E06_14) • Received Interventions for Pain Relief <u>And</u> • Values indicate a reduction in pain from the first E14_23 “pain scale” assessment to second E14_23 “pain scale” assessment 	<ul style="list-style-type: none"> • Pain Scale (E14_23) • Age (E06_14) • Age Units (E06_15) • Date of Birth (E06_16) • Date/Time Vital Taken (NEMSIS E14_01) • Date Time Medication Administered (NEMSIS E18_01) • Date Time Procedure Performed Successfully (NEMSIS E19_01)
Exclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> • Records with only one value for E14_23 or which fail a logic test; or have no value recorded for NEMSIS E14_01, or who have at least two but those values have no clear associated values for E14_01 	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	

60 • California EMS System Core Quality Measures

Benchmark Analysis	(TBD)
Rationale for Data	<input type="checkbox"/> Administration of pain treatment is important to determine the effectiveness of medicine given to patient to reduce patient pain scale level
References	NEMSIS Core Measure, Indicator 6.1

ENDOTRACHEAL INTUBATION SUCCESS RATE

MEASURE SET	Performance of Skills	
SET MEASURE ID #	SKL-1	
PERFORMANCE MEASURE NAME	Endotracheal intubation success rate	
Description	Successful intubation within two attempts.	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	All endotracheal intubation attempts	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Events in which E19_03 “procedure” has values indicating intubation such as 96.040 “endotracheal intubation” or 96.041 “airway – intubation, other (stoma, nasal)” with related element E19_05 “number of procedure attempts” 	<ul style="list-style-type: none"> Procedure (E19_03) Attempts (E19_05)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	All Successful endotracheal intubations, defined as success within 2 attempts.	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Events in which E19_03 “procedure” has values indicating intubation such as 96.040 “endotracheal intubation” or 96.041 “airway – intubation, other (stoma, nasal)” with related element E19_05 “number of procedure attempts” <p>And</p> <ul style="list-style-type: none"> E19_05 “number of procedure attempts” value listed as one or two; and E19_06 “Procedure successful” noted as value of 1 “yes” 	<ul style="list-style-type: none"> Procedure (E19_03) Attempts (E19_05) Procedure Successful (E19_06)

62 • California EMS System Core Quality Measures

Exclusion Criteria	Criteria	Data Elements
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is $N/D = \%$	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> <input type="checkbox"/>	
References	CEMSIS Core Measures	

END-TIDAL CO2 PERFORMED ON ANY ENDOTRACHEAL INTUBATION

MEASURE SET	Performance of Skills	
SET MEASURE ID #	SKL-2	
PERFORMANCE MEASURE NAME	End-tidal CO2 performed on any successful endotracheal intubation	
Description	Percentage of intubated patients where end-tidal CO2 or capnography is performed.	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	All successful endotracheal intubations	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Events in which E19_03 “procedure” has values indicating intubation such as 96.040 “endotracheal intubation” or 96.041 “airway – intubation, other (stoma, nasal)” with related element E19_05 “number of procedure attempts” E19_05 “number of procedure attempts” value listed as one or two; and E19_06 “Procedure successful” noted as value of 1 “yes” 	<ul style="list-style-type: none"> Procedure (E19_03) Attempts (E19_05) Procedure Successful (E19_06)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	All endotracheal intubations where End-Tidal CO2 measurement was performed	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> Events in which E19_03 “procedure” has values indicating intubation such as 96.040 “endotracheal intubation” or 96.041 “airway – intubation, other (stoma, nasal)” with related element E19_05 	<ul style="list-style-type: none"> Procedure (E19_03) Attempts (E19_05) Procedure Successful (E19_06)

64 • California EMS System Core Quality Measures

	<p>“number of procedure attempts”</p> <ul style="list-style-type: none"> • E19_05 “number of procedure attempts” value listed as one or two; and • E19_06 “Procedure successful” noted as value of 1 “yes” <p><u>And</u></p> <ul style="list-style-type: none"> • E19_03 “procedure” has values of 96.992 “airway-end tidal CO₂ intubation” or 89.391 “capnography” 	
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is N/D =%	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> <input type="checkbox"/>	
References	CEMSIS Core Measures	

AMBULANCE RESPONSE TIME BY AMBULANCE ZONE (EMERGENCY)

MEASURE SET	Response and Transport	
SET MEASURE ID #	RST-1	
PERFORMANCE MEASURE NAME	Ambulance response time by ambulance zone (Emergency)	
Description	90 th percentile time value of the Ambulance Response time in Ground Ambulance Transport Zone as defined by the EMS Plan	
Type of Measure	Process	
Reporting Value and Units	Time (minutes and seconds)	
Continuous Variable Statement (population)	Time (in minutes) from time ambulance is en route to arrival at the scene for emergency responses (Code 3) to patients by BLS, LALS, or ALS ambulances. The 90 th percentile time interval from “unit en route date/time” (E05-05) in an emergency to EMS “unit arrived on scene date/time” (E05-06), for a given period of time	
Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> All events in a particular ambulance zone E02_04 “type of service requested” has value 30 “911 response (scene)”; and E02_05 “Primary role of the unit” value is 75 “transport”; E02_20 “response mode to scene” is 390 “lights and sirens”; Values for E05_05 “unit en route date/time” and E05_06 “unit arrived on scene date/time” are present and logical. 	<ul style="list-style-type: none"> Ambulance Zone (Ground Ambulance Transport EOA area as defined by EMS plan) Primary role of unit (E02_05) Type of Service Requested (E02_04) Response Mode to Scene (E02_20) Unit En Route Date/Time (E05_05) Unit Arrived on Scene Date/Time (E05_06)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is the 90 th Percentile of the given numbers or distribution in their ascending order.	
Example of Final Reporting Value (number and units)	8 minutes 30 seconds	

Sampling	Yes
Aggregation	Yes
Blinded	Yes
Minimum Data Values	30
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.
Suggested Display Format & Frequency	Process control or run chart by month
Suggested Statistical Measures	90 th Percentile Measurement. Aggregate measure of central tendency and quantile (fractile) measurement to determine the span of frequency distributions.
Trending Analysis	Yes
Benchmark Analysis	(TBD)
Rationale for Data	<input type="checkbox"/> The early arrival of ambulance transport allows for the availability of those services to transport a patient from the scene in a timely manner. Common field protocols indicate that patients who have experienced severe traumatic injuries, chest pain, stroke, or other life-threatening emergencies should have a response time of less than 9 minutes. <input type="checkbox"/>
References	NEMSIS Core Measure Indicator 10.2

AMBULANCE RESPONSE TIME BY AMBULANCE ZONE (NON-EMERGENCY)

MEASURE SET	Response and Transport	
SET MEASURE ID #	RST-2	
PERFORMANCE MEASURE NAME	Ambulance response time by ambulance zone (non-emergency)	
Description	90 th percentile time value of the Ambulance Response time in Ground Ambulance Transport Zone as defined by the EMS Plan	
Type of Measure	Process	
Reporting Value and Units	Time (minutes and seconds)	
Continuous Variable Statement (population)	Time (in minutes) from time ambulance is en route to arrival at the scene for non-emergency (Code 2) responses to patients by BLS, LALS, or ALS ambulances. The 90 th percentile time interval from “unit en route date/time” (E05_05) in an emergency to EMS “unit arrived on scene date/time” (E05_06), for a given period of time	
Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> • All events in a particular ambulance zone • E02_04 “type of service requested” has value 30 “911 response (scene)”; and • E02_05 “Primary role of the unit” value is 75 “transport”; • E02_20 “response mode to scene” is 395 “no lights and sirens”; • Values for E05_05 “unit en route date/time” and E05_06 “unit arrived on scene date/time” are present and logical. 	<ul style="list-style-type: none"> • Ambulance Zone (Ground Ambulance Transport EOA area as defined by EMS plan) • Primary role of unit (E02_05) • Type of Service Requested (E02_04) • Response Mode to Scene (E02_20) • Unit En Route Date/Time (E05_05) • Unit Arrived on Scene Date/Time (E05_06)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is the 90 th Percentile of the given numbers or distribution in their ascending order.	
Example of Final Reporting Value (number and units)	8 minutes 30 seconds	

68 • California EMS System Core Quality Measures

Sampling	Yes
Aggregation	Yes
Blinded	Yes
Minimum Data Values	30
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.
Suggested Display Format & Frequency	Process control or run chart by month
Suggested Statistical Measures	90 th Percentile Measurement. Aggregate measure of central tendency and quantile (fractile) measurement to determine the span of frequency distributions.
Trending Analysis	Yes
Benchmark Analysis	(TBD)
Rationale for Data	<input type="checkbox"/> The early arrival of ambulance transport allows for the availability of those services to transport a patient from the scene in a timely manner. Common field protocols indicate that patients who have experienced severe traumatic injuries, chest pain, stroke, or other life-threatening emergencies should have a response time of less than 9 minutes. <input type="checkbox"/>
References	NEMSIS Core Measure Indicator 10.2

TRANSPORT OF PATIENTS TO HOSPITAL

MEASURE SET	Response and Transport	
SET MEASURE ID #	RST-3	
PERFORMANCE MEASURE NAME	Transport of patients to hospital	
Description	The percentage of EMS Patients transported to a General Acute Care Hospital with a Basic Permit for emergency services.	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	All 911 incidents which requested or required a response by at least one EMS unit, and the unit arrived at scene	
Denominator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> All unique EMS incidents in a particular ambulance zone E02_04 “type of service requested” has value 30 “911 response (scene)”; and E02_05 “Primary role of the unit” value is 75 “transport”; E02_20 “response mode to scene” is 3905 “lights and sirens”; Values for E05_05 “unit en route date/time” and E05_06 “unit arrived on scene date/time” are present and logical. 	<ul style="list-style-type: none"> Ambulance Zone (Ground Ambulance Transport EOA area as defined by EMS plan) Incident Number (E02_02) Primary role of unit (E02_05) Type of Service Requested (E02_04) Response Mode to Scene (E02_20) Unit En Route Date/Time (E05_05) Unit Arrived on Scene Date/Time (E05_06)
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Numerator Statement (sub-population)	All patients who received transport to a General Acute Care Hospital, with a Basic Permit, by BLS, LALS, or ALS Ambulances	
Numerator Inclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	<ul style="list-style-type: none"> All unique EMS incidents in a particular ambulance zone E02_04 “type of service requested” has value 30 “911 response (scene)”; and E02_05 “Primary role of the unit” value is 75 “transport”; E02_20 “response mode to 	<ul style="list-style-type: none"> Ambulance Zone (Ground Ambulance Transport EOA area as defined by EMS plan) Incident Number (E02_02) Primary role of unit (E02_05) Type of Service Requested (E02_04) Response Mode to Scene

70 • California EMS System Core Quality Measures

	<p>scene” is 3905 “lights and sirens”;</p> <ul style="list-style-type: none"> • Values for E05_05 “unit en route date/time” and E05_06 “unit arrived on scene date/time” are present and logical <p><u>And</u></p> <ul style="list-style-type: none"> • E20_17 has a value of 5050 “hospital” 	<p>(E02_20)</p> <ul style="list-style-type: none"> • Unit En Route Date/Time (E05_05) • Unit Arrived on Scene Date/Time (E05_06) • Patient Destination (E20_17)
Exclusion Criteria	Criteria	Data Elements
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is N/D =%	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> The early arrival of ambulance transport allows for the availability of those services to transport a patient from the scene in a timely manner. The highest risk of potential risk of harm and liability is non-transport decisions.	
References	CEMSIS Core Measures	

OUT-OF-HOSPITAL CARDIAC ARRESTS RECEIVING BYSTANDER CPR

MEASURE SET	Cardiac Arrest	
SET MEASURE ID #	PUB-1 (To Be Measured in 2014)	
PERFORMANCE MEASURE NAME	Out-of-hospital cardiac arrests receiving bystander (non-EMS Personnel/Responder) CPR	
Description	Emergency patients experiencing cardiac arrest who received CPR by bystanders (non-EMS Personnel/Responder)	
Type of Measure	Process	
Reporting Value and Units	(%) Percentage	
Denominator Statement (population)	All cardiac arrest patients	
Denominator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> Patients having a recorded E11_01 “cardiac arrest” value of 2240 “yes, Prior to EMS arrival” or value of 2245; E11_02 “cardiac arrest etiology” value of 2250 “presumed cardiac”; E11_03 “resuscitation attempted” values 2280 “attempted defibrillation” or 2285 “attempted ventilation” or 2290 “initiated chest compressions” 	<ul style="list-style-type: none"> All unique EMS incidents in a particular ambulance zone Cardiac Arrest (E11_01) Cardiac Arrest Etiology (E11_02) Resuscitation Attempted (E11_03)
Exclusion Criteria	Criteria	Data Elements
	None	
Numerator Statement (sub-population)	Patients experiencing cardiac arrest who received CPR by bystanders (non-EMS Personnel/Responder)	
Numerator Inclusion Criteria	Criteria	Data Elements
	<ul style="list-style-type: none"> Patients having a recorded E11_01 “cardiac arrest” value of 2240 “yes, Prior to EMS arrival” or value of 2245; E11_02 “cardiac arrest etiology” value of 2250 “presumed cardiac”; 	<ul style="list-style-type: none"> Cardiac Arrest (E11_01) Cardiac Arrest Etiology (E11_02) Resuscitation Attempted (E11_03) Prior Aid Performed By (E09_02)

72 • California EMS System Core Quality Measures

	<ul style="list-style-type: none"> E11_03 “resuscitation attempted” values 2280 “attempted defibrillation” or 2285 “attempted ventilation” or 2290 “initiated chest compressions” <u>And</u> E09_02 “prior aid performed by” has a value of 1205 “lay person”, and CPR was performed. 	
Exclusion Criteria	<u>Criteria</u>	<u>Data Elements</u>
	None	
Indicator Formula Numeric Expression	The formula is to divide (/) the numerator (N) by the denominator (D) and then multiply (x) by 100 to obtain the (%) value the indicator is to report. Therefore the indicator expressed numerically is N/D =%	
Example of Final Reporting Value (number and units)	90%	
Sampling	Yes	
Aggregation	Yes	
Blinded	Yes	
Minimum Data Values	30	
Data Collection Approach	<input type="checkbox"/> Retrospective data sources for required data elements include administrative data and pre-hospital care records. <input type="checkbox"/> Variation may exist in the assignment of coding; therefore, coding practices may require evaluation to ensure consistency.	
Suggested Display Format & Frequency	Process control or run chart by month	
Suggested Statistical Measures	Mean (x); Mode (m)	
Trending Analysis	Yes	
Benchmark Analysis	(TBD)	
Rationale for Data	<input type="checkbox"/> Cardiac Arrest survival has been shown to increase with early CPR and Automated External Defibrillation <input type="checkbox"/>	
References	CEMSIS Core Measures	

California EMS System Core Quality Measures

Edmund G. Brown Jr.
Governor
State of California

Diana S. Dooley
Secretary
Health and Human Services Agency

Howard Backer, MD, MPH, FACEP
Director
Emergency Medical Services Authority

EMSA Publication #166 – Appendix E
Released January 2013

www.emsa.ca.gov